



ACHIEVING SUSTAINABLE DEVELOPMENT IN LIBYA THROUGH KARST GEOCHEMISTRY



Osama R. Shaltami¹, Mohamed M. Dulayoum², Mustafa A. Ben Hkoma^{3*}, Abobakar E. Algomati⁴,
Firas Khamis Muhammed⁵, Kamal Abraheem Almahdi⁶

¹ Department of Earth Sciences, Faculty of Science, Benghazi University, Libya

² Libyan Centre for Studies & Researches in Environmental Sciences & Technology

¹ Libyan Centre for Sustainable Development Researches

⁴ Department of Engineering Geology, Faculty of Engineering, University of Bright Star, Libya

^{5,6} Department of Earth Sciences, Faculty of Science, Omar Al-Mukhtar University, Libya

³ Libyan Center for Olive Tree Research - Libyan Authority for Scientific Research

*Corresponding author: E-mail addresses: m_hkoma2017@yahoo.com

Volume: 5

Issue: 2

Page Number: 105 - 112

Keywords: Karst Geochemistry, Sustainable Development, Libya.

Copyright: © 2025 by the authors. Licensee The Derna Academy for Applied Science (DAJAS). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) License (<https://creativecommons.org/licenses/by/4.0/>).



Received: 25\11\2025

Accepted: 03\12\2025

Published: 04\12\2025

<https://doi.org/10.71147/tq2wa369>



ABSTRACT

Background: Karst in Libya is primarily found in the Al Jabal Al Akhdar region and Benghazi plain in the northeast. Karst is also observed in the Tripolitanian pre-desert region in the northwest and the Tibesti Mountains in the south. Accomplishing sustainable development in Libya through karst geochemistry involves a three-pronged approach: water resource management, hazard mitigation, and eco-tourism, all supported by ongoing scientific research. Karst systems are vital for identifying and protecting groundwater resources, which are crucial for water supply in Libya. Geochemical analysis helps monitor water quality and vulnerability to pollution, enabling better management of these resources and mitigation of hazards. Furthermore, a deeper understanding of karst systems supports the creation of geoparks and a sustainable tourism industry that conserves the environment and engages local communities. **Results:** Karst features in Libya are primarily concentrated in the Al Jabal Al Akhdar and Benghazi Plain regions, with secondary occurrences in the Tripolitanian pre-desert and Tibesti Mountains. Crucially, karst aquifers serve as vital but highly vulnerable groundwater sources, facing significant threats from pollution, overexploitation, and geomorphological hazards, necessitating urgent intervention. To address this, the study identifies a three-pronged approach for sustainable development: water resource management, hazard mitigation, and eco-tourism. Furthermore, the paper highlights that geochemical analysis is fundamental for monitoring water quality, tracing pollutant transport, and accurately delineating vulnerable zones within these complex systems. **Conclusion:** The paper concludes that karst geochemistry is essential for understanding and protecting Libya's vulnerable water resources, particularly in the critical Al Jabal Al Akhdar and Benghazi regions. To ensure sustainability, the primary recommendation is the immediate implementation of sophisticated monitoring networks and vulnerability mapping tools, alongside the development of specialized climate change adaptation strategies, all aimed at comprehensively understanding the hydrodynamic behavior and preserving the integrity of these sensitive karst ecosystems.

1. INTRODUCTION

Karst describes a landscape shaped by the dissolution of soluble bedrock like limestone, marble, and gypsum, leading to unique features. This process creates underground drainage systems where water can flow rapidly through conduits, making these aquifers highly productive but also vulnerable to contamination (e.g., Viles, 2003; Novel et al., 2007; Waele et al., 2011, 2015; Szczygieł et al., 2018; Onac and van Beynen, 2021; Zerga, 2024; Hong et al., 2026). Karst in siliceous rock is a phenomenon where landforms like caves, sinkholes, and towers form in very insoluble siliceous rocks over long periods of time, unlike classic karst that forms in soluble rocks like limestone. This "siliceous karst" is driven by processes that enhance the slow dissolution of the rock, often due to factors like acidity from organic acids in rain, specific mineral content, high rainfall, and long-term weathering (e.g., Piccini, 1995; Vaquero-Rodríguez, 2004; Vasconcelos et al., 2013; Hardt et al., 2022). Karst landscapes are ecologically and biologically important due to their role in providing freshwater habitats and aquifers for drinking water, and for supporting diverse and unique life forms adapted to both surface and subterranean environments. They are critical for biodiversity conservation because they harbor specialized species and endemic organisms, and also play a significant role in global carbon cycling (e.g., Struebig et al., 2009; Forti, 2015; Li et al., 2021; Stanković, 2023; Tarkowska-Kukuryk et al., 2025). Karst areas contain a wide variety of mineral resources, including placers of gold, tin, copper, uranium, mercury, vanadium, iron, manganese, antimony, wolframite, columbite, tantalite, bauxite, leadzinc, phosphorite, diamond, ruby, sapphire, spinel, barite, fluorite, onyx marble, clays, clay pigments, sands, coal, and peat. The fluid deposits comprise water, oil, and gas. (Filippov, 2004). Karst landscapes have significant hydrological importance due to their unique ability to store and transmit large amounts of groundwater, which serves as a vital source of freshwater for a quarter of the world's population. They are also characterized by rapid water movement through underground conduits, leading to unique hydrological processes, but making them highly vulnerable to pollution because of the lack of natural filtering. Karst hydrology is crucial for both providing drinking water and for understanding complex water systems, though it presents challenges for water management due to its sensitivity (e.g., White, 2002; Jukić and Denić-Jukić, 2006; Long, 2009; Naughton et al., 2012; Lipar and Ferk, 2015; Braitenberg et al., 2018; Bartolomé et al., 2021; Peng et al., 2025). Karst is critically important for oil and gas because it forms significant hydrocarbon reservoirs, with up to 50% of global reserves found in carbonate rock, often within karst systems. The dissolution and fracturing in karst landscapes create porous and permeable spaces where oil and gas accumulate, and understanding karstification is essential for exploration and development (e.g., Chaojun et al., 2010; He et al., 2013; Dai et al., 2017; Almalikee et al., 2020; Hao et al., 2023; Wang et al., 2025). Karst's cultural importance stems from its historical role as a human habitat and resource, with deep connections to spirituality, shelter, and traditional practices. Today, karst landscapes are valued for cultural heritage, scientific research, and economic development through geotourism (e.g., Day, 2010; Duli et al., 2019; Khalaf, 2022). Karst environments are scientifically and educationally important because they provide a unique window into Earth's history through cave deposits and morphology, serve as natural laboratories for diverse fields like biology, geology, and hydrology, and are vital for understanding climate change. Educationally, they are used as natural classrooms for multidisciplinary learning and as a tool for local economic development through geotourism and conservation efforts (e.g., Guo and Jiang, 2011; Forti, 2015; Sulistiyowati et al., 2021). Karst geochemistry studies the chemical processes that occur when water interacts with soluble rocks, creating unique landscapes (e.g., Wang et al., 2006; Han et al., 2010; Mongelli et al., 2014; D'Angeli et al., 2017; Sracek et al., 2019; Abedini et al., 2022). It is crucial for understanding how these systems form and evolve, and for assessing the impact of pollution and climate change. A number of parameters, including the Mg/Ca ratio, can be utilized to determine historical hydrological conditions in speleothems (e.g., Cruz et al., 2007; Duan et al., 2012). The temperature-dependent Mg partitioning into speleothems is usually outweighed by the notable fluctuations in the drip-water Mg/Ca ratio (e.g., Fairchild and Treble, 2009; Duan et al., 2024). The seasonal variations in the Mg/Ca ratio are thought to be exclusively explained by temperature (e.g., Carlson et al., 2018). The Mg/Ca ratio in speleothems can also be a reasonable paleotemperature parameter, but its interpretation is complex and depends on specific cave conditions, such as stable source water chemistry and a strong temperature sensitivity of magnesium partitioning. While traditionally used for hydrological data, some recent studies have shown a strong correlation between Mg/Ca ratios and cave air temperature, particularly when source water Mg/Ca is relatively constant. Major ion chemistry, trace elements, and rare earth elements (REEs) are used to study karst aquifers by providing distinct signatures that reveal how water moves, mixes, and interacts with rocks. Major ions define the overall water chemistry, trace elements identify specific sources, and REEs offer unique patterns that act as natural tracers to reveal complex processes that simpler methods miss. This combined approach allows researchers to create accurate conceptual models of the aquifer system, assess its vulnerability, and manage its water resources more effectively (e.g., Tran et al., 2023).

Adsorbed silica in stalagmites serves as a climate proxy, with a higher Si/Ca ratio generally indicating drier conditions due to the suppression of calcium carbonate precipitation and a concentration of silica in the remaining drip water. The Si/Ca ratio in speleothems is influenced by several factors, including wind-blown silicate supply, soil weathering, rainfall dilution, and the precipitation rate of calcite. A direct correlation between high Si/Ca and high stable oxygen isotopes ($\delta^{18}O$) in a stalagmite can strengthen the interpretation that regional rainfall was a major control on the speleothem's silica content (e.g., Hu et al., 2005).

2. STATEMENT OF THE PROBLEM

The study's problem lies in the acute threat to the sustainability of vital groundwater resources housed within Libya's karst systems, particularly in critical areas like the Al Jabal Al Akhdar and the Benghazi plain. These systems are crucial water sources but are characterized by high vulnerability to pollution, overexploitation, and geomorphological hazards. Despite the recognized importance of these resources and the potential role of karst geochemistry in providing the necessary understanding for effective management and protection, there is a significant knowledge and implementation gap regarding the lack of advanced monitoring networks, systematic vulnerability mapping, and planned climate change adaptation strategies. This deficit severely limits Libya's ability to achieve sustainable development and fully leverage the economic potential (such as eco-tourism) inherent in these unique geological areas.

3. LITERATURE REVIEW

Karst landscapes are prominent in northeast Libya, particularly in the Al Jabal Al Akhdar region and Benghazi plain (e.g., El Amawy et al., 2010; AlShible, 2019; Abousaif et al., 2022; Zafir et al., 2025), where limestone bedrock has dissolved to form extensive networks of caves (Fig. 1a), solution cavities (Fig. 1b), collapse dolines (Fig. 1c), karren (Fig. 1d), and springs (Fig. 1e). These formations are shaped by lithology, structural tectonics, and hydrology, and include significant sites like the Ayn Zayanah system. The northwestern part of Libya is characterized by significant karst landforms, especially in the Tripolitanian pre-desert region, and includes diverse features like caves and dolines (e.g., Hunt et al., 1985). Karst in the south, where the Tibesti Mountains are located, is less detailed in available sources, but the area is generally characterized by a mountainous, arid landscape with limited hydrological information. However, many of these unique karst landscapes are now under threat from rapid urban development.



Fig. 1: Karst features in the Al Jabal Al Akhdar region ((a) Cave, (b) solution cavities, (c) collapse doline, and (d) karren) and Benghazi plain ((e) Ayn Al Majdoub Lake).

4. THE ROLE OF KARST GEOCHEMISTRY IN PROMOTING SUSTAINABLE DEVELOPMENT IN LIBYA

Libya is focusing on sustainable development through various initiatives, including the establishment of the National Development Agency (NDA) to drive strategic projects and the United Nations Development Programme (UNDP)'s partnership with the government to advance the Sustainable Development Goals (SDGs). Key efforts include improving infrastructure, supporting economic diversification beyond oil, managing water resources through the National Water Security Strategy, and developing the renewable energy sector, especially solar and wind power. Initiatives like the "Green Libya" project also aim to enhance vegetation and combat desertification.

The following considerations highlight the importance of karst geochemistry for sustainable development in Libya:

- (1) **Water source:** Karst aquifers in Libya are a critical source of water for drinking and agriculture, although they are vulnerable to pollution and overexploitation.
- (2) **Geomorphological and speleogenetic understanding:** Geochemical and geomorphological studies, like those conducted in Al Jabal Al Akhdar, are essential to understand how karst landscapes and caves have formed over time in response to tectonic phases and climate change.

(3) Hydrological behavior: Studying the geochemistry helps understand the flow of water through these systems, including lag times between rainfall and discharge, which is vital for sustainable management.

The challenges facing karst geochemistry in attaining sustainable development in Libya are as follows:

(1) Water quality and quantity: Karst aquifers are highly susceptible to contamination and fluctuations in water supply due to their permeable nature.

(2) Geomorphic hazards: The karst landscape can be prone to geomorphic hazards, and human activities can disrupt its stability.

(3) Ecosystem sensitivity: Karst systems host unique and sensitive ecosystems, both above ground and in caves. Activities like rerouting waterways can have significant negative impacts.

(4) Climate change: Climate change poses a threat to these aquifers, and their unique characteristics require specialized adaptation strategies.

(5) Lack of comprehensive monitoring: Many karst regions in Libya lack the robust monitoring networks needed to adequately manage water resources and prevent pollution.

The following are the pathways that karst geochemistry should promote sustainable development in Libya:

(1) Advanced monitoring: Implementing sophisticated monitoring equipment for both water quantity and quality is crucial for managing dynamic karst systems, as is the case in other Mediterranean karst aquifers.

(2) Improved management: Sustainable use requires more frequent observation of discharge and water quality and better-enlarged monitoring networks than are currently in place.

(3) Vulnerability mapping: Using tools like APLIS (altitude, pendiete/slope, lithology, infiltration, and soils) and COP (concentration of flow, overlaying layers, and precipitation) for vulnerability mapping, along with isotope analysis, can help identify areas at risk and inform protection strategies.

(4) Climate adaptation: Strategies must be developed to mitigate the negative effects of climate change on karst aquifers, including adaptation strategies.

(5) Integrated research: Continued research is necessary to build a comprehensive understanding of karst evolution and groundwater resources for better management.

5. CONCLUSIONS

Karst geochemistry is crucial for understanding vulnerable water resources and is vital for sustainable development in Libya, particularly in regions like Al-Jabal Al-Akhdar and Benghazi plain. Challenges include managing water quality and quantity in these highly permeable aquifers, addressing geomorphic hazards, and ensuring development doesn't negatively impact sensitive karst ecosystems and caves. Sustainable development requires implementing sophisticated monitoring, understanding the hydrodynamic behavior of karst systems, and adapting to climate change impacts.

6. References

Abedini, A., Mongelli, G. and Khosravi, M. (2022): Geochemistry of the early Jurassic Soleiman Kandi karst bauxite deposit, Irano–Himalayan belt, NW Iran: Constraints on bauxite genesis and the distribution of critical raw materials. *Journal of Geochemical Exploration*; 241: 107056.

Abousaif, A., Abuharara, A., Bousayf, A. and Atia, M. (2022): Geological and geomorphic assessment of Wadi Al-Battum, Al Jabal Al Akhdar, Libya. *Alqalam Journal of Medical and Applied Sciences (AJMAS)*; 5(1): 177-184.

Almalikee, H.S., Almayyahi, H.K. and Al-Jafar, M.K. (2020): Karst feature in Mishrif Reservoir and effect on drilling and production in Zubair oil field, Southern Iraq. *Journal of Petroleum Research and Studies (JPRS)*; 29(12): E19-E32.

AlShible, M.S. (2019): Geology of Ayn Al Majdoub karstic lake, Benghazi, NE Libya. *Libyan Journal of Science and Technology (LJST)*; 9(1): 93-97.

Bartolomé, M., Sancho, C., Benito, G., Medialdea, A., Calle, M., Moreno, A., Leunda, M., Luetscher, M., Muñoz, A., Bastida, J., Cheng, H. and Edwards, R.L. (2021): Effects of glaciation on karst hydrology and sedimentology during the Last Glacial Cycle: The case of Granito cave, Central Pyrenees (Spain). *CATENA*; 206: 105252.

- Braitenberg, C., Pivetta, T., Rossi, G., Ventura, P. and Betic, A. (2018): Karst caves and hydrology between geodesy and archeology: Field trip notes. *Geodesy and Geodynamics*; 9(3): 262-269.
- Carlson, P.E., Miller, N.R., Banner, J.L., Breecker, D.O. and Casteel, R.C. (2018): The potential of near-entrance stalagmites as high-resolution terrestrial paleoclimate proxies: Application of isotope and trace-element geochemistry to seasonally-resolved chronology. *Geochimica et Cosmochimica Acta*; 235: 55-75.
- Chaojun, Z., Chengzao, J., Benliang, L., Xiuyu, L. and Yunxiang, L. (2010): Ancient karsts and hydrocarbon accumulation in the middle and western parts of the North Tarim uplift, NW China. *Petroleum Exploration and Development*; 37(3): 263-269.
- Cruz Jr., F.W., Burns, S.J., Jercinovic, M., Karmann, I., Sharp, W. D. and Vuille, M. (2007): Evidence of rainfall variations in Southern Brazil from trace element ratios (Mg/Ca and Sr/Ca) in a Late Pleistocene stalagmite. *Geochimica et Cosmochimica Acta*; 71(9): 2250-2263.
- Dai, X., Zhang, M., Jiang, Q. and Feng, Z. (2017): Karst reservoirs seismic prediction of Lower Permian Maokou Formation in central Sichuan Basin, SW China. *Petroleum Exploration and Development*; 44(1): 79-88.
- D'Angeli, I.M., Serrazanetti, D.I., Montanari, C., Vannini, L., Gardini, F. and DeWaele, J. (2017): Geochemistry and microbial diversity of cave waters in the gypsum karst aquifers of Emilia Romagna region, Italy. *Science of the Total Environment*; 598: 538-552.
- Day, M. (2010): Human interaction with Caribbean karst landscapes: Past, present and future. *ACTA CARSOLOGICA*; 39(1): 137-146.
- Duan, W., Cai, B., Tan, M., Liu, H. and Zhang, Y. (2012): The growth mechanism of the aragonitic stalagmite laminae from Yunnan Xianren Cave, SW China revealed by cave monitoring. *Boreas*; 41(1): 113-123.
- Duan, W., Wu, S., Wang, X., Shen, J., Xiao, Z., Cui, L. and Tan, M. (2024): Ma/Ca Ratio in stalagmites as a potential palaeo- temperature indicator. *Geophysical Research Letters*; 51: e2024GL108498.
- Duli, A., Mulyadi, Y. and Rosmawati (2019): The mapping out of Maros-Pangkep karst forest as a cultural heritage conservation. The 1st Biennial Conference on Tropical Biodiversity, IOP Conference Series: Earth and Environmental Science; 270: 012014.
- El Amawy, M.A., Muftah A.M. and Abdelmalik, M.B. (2010): Karst development and structural relationship in the Tertiary rocks of the western part of Al Jabal Al Akhdar, NE Libya: A case study in Qasr Libya area. 3rd International Symposium Karst Evolution in the South Mediterranean Area, *Speleologia Iblea, Ragusa, Italy*; 14: 173-189.
- Fairchild, I.J. and Treble, P.C. (2009): Trace elements in speleothems as recorders of environmental change. *Quaternary Science Reviews*; 28(5-6): 449-468.
- Filippov, A.G. (2004): Mineral deposits in karst. In: Gunn, J. (Eds.), *Encyclopedia of Caves and Karst Science*, Taylor and Francis; pp. 514-515.
- Forti, P. (2015): The scientific and socio-economic importance of karst and caves and their vulnerability. Brief for Global Sustainable Development Report (GSDR); pp. 1-3.
- Guo, F. and Jiang, G. (2011): Karst groundwater management through science and education. *Open Journal of Geology*; 1: 45-50.
- Han, G., Tang, Y. and Xu, Z. (2010): Fluvial geochemistry of rivers draining karst terrain in Southwest China. *Journal of Asian Earth Sciences*; 38(1-2): 65-75.
- Hao, Y., Bell, R., Rao, Y., Fan, R. and Wang, Y. (2023): Prediction of Permian karst reservoirs in the Yuanba gas field, northern Sichuan Basin, China. *Marine and Petroleum Geology*; 154: 106319.
- Hardt, R., Rodet, J. and Zogbi, L.D. (2022): Sandstone and karst development in Brazil. 18th International Congress of Speleology - Savoie Mont Blanc, *Karstologia Memoires n°23 - Symposium 11 – Pseudokarst*; 3: 239-242.

- He, J., Fang, S., Hou, F., Yan, R., Zhao, Z., Yao, J., Tang, X. and Wu, G. (2013): Vertical zonation of weathered crust ancient karst and reservoir evaluation and prediction—A case study of M–M sub-members of Majiagou Formation in gas fields, central Ordos Basin, NW China. *Petroleum Exploration and Development*; 40(5): 572-581.
- Hong, Y., Li, Z., Xin, H., Pan, F., Xie, H. and Duan, L. (2026): Effective discharge for sediment transport and its geomorphological significance in karst catchments. *Journal of Hydrology*; 664(A): 134412.
- Hu, C., Huang, J., Fang, N., Xie, S., Henderson, G.M. and Cai, Y. (2005): Adsorbed silica in stalagmite carbonate and its relationship to past rainfall. *Geochimica et Cosmochimica Acta*; 69(9): 2285-2292.
- Hunt, C.O., Gale, S.J. and Gilbertson, D.D. (1985): The UNESCO Libyan Valleys Survey IX: Anhydrite and limestone karst in the Tripolitanian pre-desert. *Libyan Studies*; 16: 1-13.
- Jukić, D. and Denić-Jukić, V. (2006): Nonlinear kernel functions for karst aquifers. *Journal of Hydrology*; 328(1-2): 360-374.
- Khalaf, E.A. (2022): Karst heritage as a tourist attraction: A case study in the White Desert National Park, Western Desert, Egypt. *Geoheritage*; 14: 94.
- Li, S., Liu, C., Chen, J. and Wang, S. (2021): Karst ecosystem and environment: Characteristics, evolution processes, and sustainable development. *Agriculture, Ecosystems and Environment*; 306: 107173.
- Lipar, M. and Ferk, M. (2015): Karst pocket valleys and their implications on Pliocene–Quaternary hydrology and climate: Examples from the Nullarbor Plain, southern Australia. *Earth-Science Reviews*; 150: 1-13.
- Long, A.J. (2009): Hydrograph separation for karst watersheds using a two-domain rainfall–discharge model. *Journal of Hydrology*; 364(3-4): 249-256.
- Mongelli, G., Boni, M., Buccione, R. and Sinisi, R. (2014): Geochemistry of the Apulian karst bauxites (southern Italy): Chemical fractionation and parental affinities. *Ore Geology Reviews*; 63: 9-21.
- Naughton, O., Johnston, P.M. and Gill, L.W. (2012): Groundwater flooding in Irish karst: The hydrological characterisation of ephemeral lakes (turloughs). *Journal of Hydrology*; 470-471; 82-97.
- Novel, J., Dimadi, A., Zervopoulou, A. and Bakalowicz, M. (2007): The Aggitis karst system, Eastern Macedonia, Greece: Hydrologic functioning and development of the karst structure. *Journal of Hydrology*; 334(3-4): 477-492.
- Onac, B.P. and van Beynen, P. (2021): Caves and karst. 2nd edition, *Encyclopedia of Geology*; pp. 495-509.
- Peng, G., Gao, X., Naseem, A., Zhang, Y., Wang, X., Fu, W., Yu, F., Ma, S., Shi, W., Yi, L., Li, C., Liu, W. and Kong, S. (2025): Karst water quality, source of pollution, and health risk assessment in China. *Science of the Total Environment*; 973: 179120.
- Piccini, L. (1995): Karst in siliceous rock: karst landforms and caves in the Auyan-tepui (Est. Bolivar, Venezuela). *International Journal of Speleology*; 24(1-4): 41-54.
- Sracek, O., Geršl, M., Faimon, J. and Bábek, O. (2019): The geochemistry and origin of fluids in the carbonate structure of the Hranice Karst with the world's deepest flooded cave of the Hranická Abyss, Czech Republic. *Applied Geochemistry*; 100: 203-212.
- Stanković, V.M. (2023): Biodiversity in karst landscapes: Introduction to the special issue. *Diversity*; 15(3): 408.
- Struebig, M.J., Kingston, T., Zubaid, A., Le Comber, S.C., Mohd-Adnan, A., Turner, A., Kelly, J., Božek, M. and Rossiter, S.J. (2009): Conservation importance of limestone karst outcrops for palaeotropical bats in a fragmented landscape. *Biological Conservation*; 142(10): 2089-2096.
- Sulistiyowati, E., Setiadi, and Haryono, E. (2021): Karst and conservation research in Indonesia and its implication to education. *Young Scholar Symposium on Science Education and Environment (YSSSEE)*, *Journal of Physics: Conference Series*; 1796: 012071.
- Szczygieł, J., Golicz, M., Hercman, H. and Lynch, E. (2018): Geological constraints on cave development in the plateau-gorge karst of South China (Wulong, Chongqing). *Geomorphology*; 304: 50-63.

- Tarkowska-Kukuryk, M., Ferencz, B., Dawidek, J. and Ziólek, A. (2025): Ecological health assessment of karstic springs based on the biological integrity of benthic fauna. *Global Ecology and Conservation*; 63: e03907.
- Tran, D.A., Goeppert, N. and Goldscheider, N. (2023): Use of major ion chemistry and trace and rare earth elements to characterize hydraulic relations, mixing processes and water–rock interaction in the Dong Van karst aquifer system, Northern Vietnam. *Hydrogeology Journal*; 31: 1735-1753.
- Vaqueiro-Rodríguez, M. (2004): Karst in siliceous rocks: Pseudokarst landforms and caves in the quartz dyke of Pico Sacro (Boqueixon, Galicia). *Coruña*; 29: 247-264.
- Vasconcelos, A.M.C., de Souza, F.C.R., Rodet, J., de Oliveira, C.V. and Salgado, A.A.R. (2013): Karst developed in siliciclastic rocks at Serra Do Espinhaço Meridional, Minas Gerais (Brazil). *Karst and Caves in Other Rocks, Pseudokarst – Poster, 16th International Congress of Speleology (ICS) Proceedings, Czech Speleological Society*; pp. 326-331.
- Viles, H.A. (2003): Conceptual modeling of the impacts of climate change on karst geomorphology in the UK and Ireland. *Journal for Nature Conservation*; 11(1): 59-66.
- Waele, J.D., Gutiérrez, F. and Audra, P. (2015): Karst geomorphology: from hydrological functioning to palaeoenvironmental reconstructions. *Geomorphology*; 229: 1-2.
- Waele, J.D., Gutiérrez, F., Parise, M. and Plan, L. (2011): Geomorphology and natural hazards in karst areas: A review. *Geomorphology*; 134(1-2): 1-8.
- Wang, Y., Guo, Q., Su, C. and Ma, T. (2006): Strontium isotope characterization and major ion geochemistry of karst water flow, Shentou, northern China. *Journal of Hydrology*; 328(3-4): 592-603.
- Wang, Y., Chen, S., Wu, J., Qian, Y. and Hou, X. (2025): Karst hydrocarbon reservoir characterization and sweet spot prediction in the western Tazhong Oilfield, China. *International Journal of Energy Research*; 2025(1): 4038164.
- White, W.B. (2002): Karst hydrology: Recent developments and open questions. *Engineering Geology*; 65(2-3): 85-105.
- Zafir, A., Hassan, S., Khamees, F. and Mustafa, S. (2025): Influence of structure and stratigraphy on karst development in Ayn Hufrah, NE Libya. *International Scientific Conference Man and Karst, Speleologia Iblea, Ragusa, Italy*; XIX: 57-61.
- Zerga, B. (2024): Karst topography: Formation, processes, characteristics, landforms, degradation and restoration: A systematic review. *Watershed Ecology and the Environment*; 6: 252-269.